



Beginner's Guide to GOES-R Series Data

How to acquire, analyze, and visualize GOES-R Series data

Resources compiled by GEO Program Science

Last Updated on September 8, 2025



Document Change Record

Version	Date	Slides Affected	Description
1.0	10/21/2020	All	Initial
1.1	2/8/2021	8, 12, 15-17, 19-21, 24, 26-29	The following content was added: additional GOES-R imagery viewing platforms; AWS "S3 Explorer" user interface for file browsing; Geo2Grid software; Python packages (satpy, GOES-2-Go, goespy); section on GIS processing (converting netCDF files to GeoTIFFs, transforming data to shapefiles).
1.2	5/23/2024	5-9, 12, 14-16, 31-33	Updates include: New figures for GOES-R series satellite and GOES satellite constellation; adjustments needed post GOES-18 designation as GOES-West; updates to multiple links
1.3	9/8/2025	1, 5-9, 12-17, 19-24, 27, 28, 30-37	Updates to descriptions and figures to reflect GOES-19 as the new operational GOES-East; updates to multiple links and their descriptions; remove OCC data access description



Contents

Part 1: The GOES-R Series

- Introduction to the GOES-R Series 5
- GOES-R Series Instruments 6
- Advanced Baseline Imager (ABI) 7
- ABI Mesoscale Domains 8
- GOES-R Data Products 9
- Product Maturity and Data Availability 10

Part 2: Where Can I Access the Data?

- View GOES-R Imagery 12
- View GOES-R Imagery with AWIPS 13
- Access Data Files: NOAA CLASS 14
- Access Data Files: Amazon, Microsoft 15
- Access Data Files: Google Cloud 16
- Use Python to Retrieve Data from AWS 17

Part 3: How Can I Display the Data?

- Visualize with Command Line: Geo2Grid 19
- Visualize with Python: Basics 20
- Visualize with Python: Custom Packages 21

- Visualize with Earth Engine 22
- Technique: Radiance to Reflectance 23
- Technique: Generating Composites 24

Part 4: How Can I Process the Data Using GIS?

- Make GOES-R Series Data GIS-compatible 26
- Convert NetCDF files to GeoTIFFs 27
- Transform Discrete Data to Shapefiles 28

Part 5: Frequently Asked Questions

- How are GOES-R Series files formatted? 30
- How do I pre-process GOES-R Series data? 31
- How are AB1 L1b products georeferenced? 32
- Why is data not available on a given date? 33
- Other Questions and Contact 34

Appendix A: Acronym List 35

Appendix B: Deprecated Script Examples 36

Part 1.



The GOES-R Series





Introduction to the GOES-R Series

- Geostationary Operational Environmental Satellites (GOES) are developed, launched and operated in a collaborative effort by NOAA and NASA, and have been in operation since 1975.
- The latest generation of geostationary satellites is the GOES-R Series, with its first launch in 2016.
- The GOES-R Series is a four satellite program, which maintains two operational satellites at all times, a third standby satellite in “storage mode” on-orbit as a ready spare, and the fourth also in “storage mode”.




Figure 1. GOES-R series fleet locations as of June 2025. The geographic ranges of GOES-East and GOES-West together cover the North American continent.

Satellite	GOES-R (GOES-16)	GOES-S (GOES-17)	GOES-T (GOES-18)	GOES-U (GOES-19)
Launch date	Nov. 19, 2016	Mar. 1, 2018	Mar. 1, 2022	Jun. 25, 2024
Current status	Standby (104.7° W)	Storage (89.5° W)	Operational (137° W)	Operational (75.2° W)
Operational position	GOES-East	GOES-West	GOES-West	GOES-East
Operations start	Dec. 18, 2017	Feb. 12, 2019	Jan. 4, 2023	Apr. 7, 2025
Operations end	Apr. 7, 2025	Jan. 4, 2023	-	-

Table 1. Status summary of each GOES-R series satellite.

GOES-R Series Instruments

Earth-pointing:

- [Advanced Baseline Imager](#) (ABI) - the primary instrument for imaging Earth's weather, oceans and environment. See [Slide 7](#) for more details.
- [Geostationary Lightning Mapper](#) (GLM) - a single-channel, near-infrared optical transient detector that can identify momentary changes in an optical scene, indicating the presence of lightning.

Sun-pointing:

- [Extreme Ultraviolet and X-ray Irradiance Sensors](#) (EXIS) - monitors solar irradiance in the upper atmosphere using two primary sensors: the Extreme Ultraviolet Sensor (EUVS) and the X-Ray Sensor (XRS).
- [Solar Ultraviolet Imager](#) (SUVI) - a telescope that monitors the sun in the extreme ultraviolet wavelength range, detecting solar flares and solar eruptions, and compiling full disk solar images.
- [Compact Coronagraph](#) (CCOR) - instrument carried only on GOES-19 that images the outer layer of the sun's atmosphere (corona), helping to detect and characterize Coronal Mass Ejections (CMEs).

In-situ:

- [Magnetometer](#) (MAG for GOES-16/GOES-17, GMAG for GOES-18/GOES-19) - measures the space environment magnetic field that controls charged particle dynamics in the outer region of the magnetosphere.
- [Space Environment In-Situ Suite](#) (SEISS) - monitors proton, electron, and heavy ion fluxes in the magnetosphere using four sensors: the Energetic Heavy Ion Sensor (EHIS), the High and Low Magnetospheric Particle Sensors (MPS-HI and MPS-LO), and the Solar and Galactic Proton Sensor (SGPS).



Figure 2. GOES-R Series satellites are composed of 6 (7 on GOES-19) instruments, and powered by a solar panel array.



Advanced Baseline Imager (ABI)

- [ABI](#) is a multi-channel passive imaging radiometer that images Earth's weather, oceans and environment with 16 spectral bands (2 visible, 4 near-infrared, and 10 infrared channels).
 - [ABI Bands Technical Summary Chart](#)
- Spatial resolution is 0.5, 1, or 2 km, depending on the band (see above-linked chart).
- Geographic coverage
 - Full Disk: a circular image depicting nearly full coverage of the Western hemisphere ([GOES-East/GOES-West](#)).
 - CONUS/PACUS : a 3,000 (lat) by 5,000 (lon) km rectangular image depicting the Continental US ([CONUS](#)) (GOES-East) or the Pacific Ocean including Hawaii ([PACUS](#)) (GOES-West).
 - Mesoscale: a 1,000 by 1,000 km rectangular image. GOES-East and GOES-West both alternate between two different mesoscale geographic regions (domains). See [Slide 8](#) for a complete description of mesoscale domains.
- ABI has multiple [scan modes](#).
 - Mode 6 - The default operational mode for GOES-East & West. Produces a full disk every 10 minutes, a CONUS/PACUS image every 5 minutes, and images of both mesoscale domains every 60 seconds.
 - Mode 4 - A contingency mode. Produces only a full disk image every 5 minutes.
 - Mode 3 - The default operational mode for GOES-R satellites until April 2019 when it was replaced by Mode 6. Produced a full disk every 15 minutes, a CONUS image every 5 minutes, and images of both mesoscale domains every 60 seconds.
 - A “cooling timeline” version of Mode 3 was also utilized by GOES-17, starting in February 2020, to mitigate image saturation due to the [loop heat pipe \(LHP\) anomaly](#) during the seasons of extra solar heating (late Feb-early April, late Aug-early Oct). Mode 3 cooling timeline produces a full disk image every 15 minutes, and images of both mesoscale domains every 2 minutes.

ABI Mesoscale Domains

- Mesoscale domains are 1,000 by 1,000 km movable rectangular regions. Mesoscale scans frequently revisit an area of interest to monitor regional conditions.
- GOES-East and GOES-West each have two default domains (see Fig. 4), however the domains can be positioned anywhere within the full disk upon request. Requests are placed by National Weather Service (NWS) Weather Forecast Offices (WFOs).
 - Figure 3 shows the mesoscale requests by month.
 - Check out the [live feed of current mesoscale locations](#).
- In Mode 6, the ABI on both GOES-East and GOES-West scans either the same mesoscale domain every 30 seconds, or two separate mesoscale domains every 60 seconds.

Figure 4a. The GOES-East default mesoscale scan sectors are the East Coast and Midwest.





Figure 4b. The GOES-West default mesoscale scan sectors are the West Coast and Alaska.



GOES-R Series Mesoscale Domain Requests by Type




Figure 3. The frequency of mesoscale domain requests varies over time, peaking during summer months.



GOES-R Data Products

[Here is a list of all the GOES-R products](#) made available to the user community by NOAA.

- Level 0 (L0):
 - L0 products are observation data received directly from the satellite instruments. The data are not easily accessible nor meaningful to most users prior to processing by the ground system.
- Level 1b (L1b):
 - L1b products are calibrated and, where applicable, geographically-corrected, L0 data. This means that the data have been processed so that the values are in standard units of physical quantities.
 - For ABI, the L1b product is radiances. This is useful for users who require radiance units, instead of reflectance/brightness temperature (Kelvin) units.
 - All of the instruments have L1b products distributed except GLM, which is only provided as an L2+ product.
 - For technical information, see the Product User Guide (PUG) Volume 3: L1b Products, found on the [OSPO GOES-R Documents page](#).
- Level 2+ Products:
 - L2+ products contain environmental physical qualities, such as cloud top height or land surface temperature. Aside from the [GLM Lightning Detection Product](#), the data source for these products is the ABI L1b data.
 - The mission-critical ABI product is [Cloud and Moisture Imagery](#) (CMI), which utilizes all 16 ABI spectral bands, and is used to generate an array of products aiding forecasters in monitoring and predicting weather hazards.
 - For technical information, see the PUG Volume 5: L2+ Products, found on the [OSPO GOES-R Documents page](#).



Product Maturity and Data Availability

- Before GOES-R satellites take off, pre-launch verification determines that systems are functioning.
- However, most instrument, L1b, and L2+ product validation is fully realized after launch with post-launch product tests that use actual earth, solar, and space observations. Calibration and characterization also continue after launch to maintain data product quality.
- Product Maturity Levels, *summarized*:
 - **Beta:** Data are preliminary and non-operational; undergoing testing and initial calibration and validation. Beta products have been minimally validated and may still contain significant errors.
 - **Provisional:** Data are ready for operations. Performance has been tested and documented over a subset of conditions, locations, and periods. Validation is still ongoing; known anomalies are documented and available to the user community.
 - **Full:** Product is operational. All known anomalies are documented and shared with the user community. Performance has been tested and documented over a wide range of conditions.
- Data Availability:
 - CLASS and NCEI (see [Slide 14](#)) have all data from launch onward available to end-users.
 - Cloud-based platforms (see [Slides 15 and 16](#)) typically have only provisional data and onwards available to users. However, the length of time of the data archive may vary between vendors.
 - To determine when a particular product advanced to a certain maturity level, explore the [Peer/Stakeholder Product Validation Reviews](#).

Part 2.



Where Can I Access the Data?





View GOES-R Imagery

- There are a variety of platforms designed for viewing and downloading recent GOES-R images and animated image loops, but not for analyzing the data.
- [GOES Image Viewer](#) shows ABI bands, band composites, and GLM data from the past 20 hours.
 - For any active storm, a console describes the weather event and provides a real-time animation.
- [CIRA “RAMMB” Slider](#) provides near real-time ABI bands, composites and some L2+ products, including GLM, along with imagery from other satellites (Himawari, Meteosat, JPSS, GEO-KOMPSAT) from the past 7 days.
- [CSSP GeoSphere](#) displays only ABI data, but with a two week rolling archive. Data is processed using the open-source software Geo2Grid, which can be used by anyone (see [Slide 19](#)).
- [NASA Worldview](#) can overlay select ABI data (Channels 2 and 13, Fire Temperature, and Air Mass, Dust, and GeoColor composites) from the past 90 days with hundreds of atmospheric, oceanic, and environmental layers.
- [RealEarth](#) is another platform for intercomparison of ABI imagery with various global satellite products. ABI imagery is provided in near real-time, but only archived for three days.
- NOAA’s Space Weather Prediction Center (SWPC) provides near real-time [SUVI](#) and [CCOR](#) imagery.
 - SWPC also provides [near real-time observations](#) for the other non-imaging GOES-R space weather instruments
- Other viewing platforms:
 - [NASA SPoRT](#) illustrates ABI bands and several composites, but only has a one day archive.
 - [SSEC Viewer](#) provides near real time imagery of all ABI channels and a True Color composite.
 - For additional viewing platforms, see the [“GOES ABI \(Advanced Baseline Imager\) Realtime Imagery” web page](#).

View GOES-R Imagery with AWIPS

- AWIPS (Advanced Weather Interactive Processing System) is the software used by the National Weather Service (NWS) to display and analyze meteorological data.
- Unidata developed and supports a [modified non-operational version of AWIPS](#) which is a free and open-source software that any user can download to view GOES-R Series data through a similar lens as weather forecasting offices (WFOs).
 - CAVE (Common AWIPS Visualization Environment) is a Java application which runs on Linux, Mac, and Windows. Follow [these steps to download CAVE to your device](#).
 - AWIPS data can be accessed through the cloud using the [EDEX Server](#). When prompted by the Connectivity Preferences Dialog, select the EDEX-cloud server.
 - Read Unidata's [CAVE User Manual](#) to learn how to interact with and customize the AWIPS tools. See the [GOES East/West Section](#) for instructions specific to GOES-R channels, products, and RGB Composites.
 - For more detail, [review the entire user manual](#), with particular attention to Section 2.2.
 - See p.56-63 for image display and time cadence options, and p.53-55 for viewing preferences.




Figure 5. The AWIPS display can overlay L2+ products and visible imagery, such as Land Surface Temperature and Channel 2.



Access Data Files: NOAA CLASS

- The NOAA [Comprehensive Large Array-data Stewardship System \(CLASS\)](#) repository is the official site for accessing all available GOES-R Series Products.
 - Begin by selecting the product of interest from the search bar, such as [GOES-R Series ABI Products \(GRABIPRD\) \(partially restricted L1b and L2+ Data Products\)](#).
 - Use the Temporal and Advanced Search options to filter the data by time, date, satellite, geographic scale, and product. It is possible to include up to 10,000 files in an order. Click “Quick Search and Order”, then “Register”.
 - Fill in your contact information to create a Guest User Profile. Once you are logged into CLASS as a guest, the option to “Place Order” will appear. If you are a returning CLASS user, log in before beginning your data search.
 - When the order has been processed (up to 48 hours for large order sizes), instructions will be sent to the email address provided. The email will offer two options for accessing the file(s):
 - 1. Authenticate and download from [CLASS via FTPS](#). This requires a robust FTP client. Use “anonymous” as the FTP user ID, and your email address as the password.
 - 2. Download each file individually from the CLASS website.
- A similar but easier website to navigate is NCEI's [Archive Information Request System \(AIRS\)](#).
 - Using the AIRS system, there is no need to set up an account.
 - It is possible to filter, search, and order up to 30 days worth of files, and access up to 1,000 files per order.
 - An email will be sent with download instructions for the processed files, with options for web and FTP downloads.
 - Another interface for downloading NCEI's archive of GOES-R Series data is [NCEI Dataset Search](#).



Access Data Files: Amazon, Microsoft

[Amazon Web Service \(AWS\)](#) - ABI L1b and L2+, GLM L2, and SUVI, SEISS, MAG, and EXIS L1b products are available in AWS S3 Buckets. These open datasets can be accessed by the public from AWS for free.

- User interfaces for browsing and downloading the available products:
 - The AWS S3 Explorer interface can be used to browse the individual satellite buckets ([GOES-16](#), [GOES-17](#), [GOES-18](#), [GOES-19](#)).
 - Alternatively, users with AWS accounts can browse the buckets via the [AWS S3 Console](#).
- Options for scripted/bulk downloading:
 - [AWS Command Line Interface](#), [rclone](#) (see [rclone tutorial for GOES data access](#)), and the [Python s3fs library](#) (see [Slide 17](#) for more information on using Python to download data).
- See the [README file](#) under “Documentation” on the AWS page for GOES-specific instructions and file naming.
- ABI L1b Full Disk and CONUS data are being [reprocessed](#) and archived in the AWS buckets.
- A separate [AWS S3 Bucket holds CCOR data](#) (GOES-19 only).

[Microsoft Planetary Computer](#) - GOES-R Series products are stored in a Azure blob container.

- The products currently available are: ABI Cloud & Moisture Imagery, select ABI L2+ derived products, and GLM L2 Lightning Detection which can be accessed from Azure for free.



Access Data Files: Google Cloud

- [Google Cloud](#) - ABI L1b and L2+, GLM L2, and SUVI, SEISS, MAG, and EXIS L1b products are available in different buckets for [GOES-16](#), [GOES-17](#), [GOES-18](#), and [GOES-19](#).
 - GOES-R data can be browsed and pulled directly from the Google Cloud website. The data itself is stored in a Google Cloud Storage Bucket, while the indexed metadata is available in BigQuery.
 - The buckets can be filtered by product name and date, but be prepared with the product abbreviation and the day-of-the-year of interest, because there are few descriptors on the Google Cloud website.
 - Downloading the data requires authentication with a Google account (not Google Cloud specific).
 - Without a Google account, data access is still possible using a Cloud Storage API link.
 - BigQuery offers [1TB of querying per month](#) for free. Additionally, a [Google Cloud free trial](#) provides \$300 in credit, valid for 90 days, which can be used to explore other Google Cloud services.
 - Follow the “[How to process weather satellite data in real-time in BigQuery](#)” tutorial to visualize GOES-R data using Google Cloud and the gsutil command-line utility.
- [Google Earth Engine](#) (GEE) - A cloud-based platform for geospatial analysis. This service runs through Google Cloud, and pulls GOES-R datasets from the Google Cloud buckets.
 - Only [two GOES-R L2+ products are currently available in the GEE Data Catalog](#): Multi-band Cloud and Moisture Imagery (MCMi) and Fire Detection and Characterization (FDC).
 - [Apply for a free noncommercial GEE account](#) to use this platform. It may take 1-2 days for account approval.
 - See [Slide 22](#) for more information on how to use GEE as a tool for image analysis and visualization.



Use Python to Retrieve Data from AWS

- The following sample scripts demonstrate two different ways to retrieve GOES-R files from AWS using Python. The scripts must be edited to use the correct prefix for the file of interest.
 - For GOES-R file naming conventions, see the AWS [README](#) documentation.
- [Script #1](#): “Visualize GOES-16 Data from S3” by Hamed Alemohammad
 - This script bypasses downloading data to a local device by pulling data directly from the AWS S3 Bucket, visualizing the data, and saving the image as a png. Recommended edits to Script #1 follow, based on a 1/6/21 review:
 - In [2]: Adjust information to reflect the specific satellite/product/date of interest. Be wary that **not** all products have data for multiple bands (if not, remove “band” attribute), or have images for every date.
 - In [5]: Change “M3” to “M6” (mode 3 to mode 6) if the imagery of interest was created after April 2, 2019.
 - This reflects the switch in ABI’s default scan mode. See [Slide 7](#) for more information on scan modes.
 - In [8]: Change “Rad” to other variable name, if the product of interest is not an ABI L1b Radiance Product.
- [Script #2](#): “Download GOES AWS” by Brian Blaylock, Ph.D.
 - This simple script demonstrates how to do a scripted download of GOES-R files from AWS using anonymous credentials. Files are stored at a local directory. Recommended edits to Script #2 follow, based on a 1/6/21 review:
 - Line 12 : Adjust the bucket name from 's3://noaa-goes16/' to “goes17”, “goes18”, or “goes19” in place of “goes16” if applicable.
 - Line 17 : Correct the prefix from 'noaa-goes17/ABI-L2-MCMIPC/2019/240/00/' to the file of interest.
- The GOES-2-Go and goespy Python packages can further streamline the download (see [Slide 21](#)).

Part 3.



How Can I Display the Data?



Visualize with Command Line: Geo2Grid

- The free [CSSP Geo2Grid](#) software package is a set of command line tools which facilitate the creation of projected satellite images, developed at [CIMSS/SSEC](#).
- Installation and setup:
 - Install the Geo2Grid software package on a device running Anaconda/Miniconda:
 - In command line, run: `conda create -c conda-forge -n geo2grid polar2grid`, then `conda activate geo2grid`
 - Download ABI L1b files of interest, and store in a local directory.
 - Alternatively, use pip to install:
 - Install the PyPI package: `pip install polar2grid`
 - Or, clone the [git repository](#) and install using: `pip install .`
- Use the [ABI L1b Reader](#) to read data, and write to single band GeoTIFF files:
 - `geo2grid -r abi_l1b -w geotiff -f <path_to_files>`
- RGB composites can be generated by specifying `-p [product]` (Figure 6):
 - `geo2grid -r abi_l1b -w geotiff -p true_color -f <path_to_files>`
 - Products: True color*, natural color* air mass, ash, dust, fog, and night microphysics.
 - *These RGBs are ratio-sharpened, enhanced, and an atmospheric correction is applied.
- Other functions include: remapping data to a different projection, subsetting data to a geographic bounding box, and setting the file output format to a png.
- For more information, see [Geo2Grid Basics](#) and [Examples for Working with ABI Data](#).





Figure 6. GeoTIFFs created using Geo2Grid. 6a (top): GOES-16 CONUS true color. 6b (bottom): GOES-17 PACUS natural color.

Visualize with Python: Basics




Figure 7. GOES-16 visualized with Python. 7a (top) ABI L1b Channel 2 Full Disk Radiances from Jan 5th, 2021. 7b (bottom): True Color CONUS from June 4th, 2020.

- Import data to Python directly from AWS (see [Slide 17](#) and [Slide 19](#) for examples), or from a local directory.
- Visualize GOES-R imagery using Python packages: xarray and matplotlib.
 - Xarray processes GOES-R data as multidimensional arrays. Matplotlib plots data to generate images with legends. Figure 7a adapted from [Script #1](#) output.
- Georeference GOES-R imagery using Cartopy package.
 - Cartopy helps georeference data, so that geographic features like state borders can be overlaid. Cartopy should be used instead of Basemap, a package that is now deprecated.
 - [Script #3](#): “Mapping GOES-16 True Color” by Brian Blaylock, Ph.D. A step by step explanation for how to render and georeference a GOES-16 ABI true color image. Figure 7b created by following this script. Recommended edits to Script #3 follow, based on a 6/27/25 review:
 - Under “Open the GOES-16 NetCDF File”, “Overlay Nighttime IR when dark”, “Can we make plots for a Mesoscale scan?”, and “Can we do this for a Full Disk Scan? It’s possible...” sections, the linked files are no longer accessible. Instead, download files locally (e.g., from [AWS](#)) and reference those in the `FILE` line.
 - Under “Using other projections”, “Plot with Cartopy: Plate Carrée Cylindrical Projection”, and “Can we make plots for a Mesoscale scan?” sections, the `ax.imshow` extent lines need to reference the x and y values. Replace the `extent` lines with `extent=(x.values.min(), x.values.max(), y.values.min(), y.values.max())`.
- [Script #4](#): “Accessing GOES data with the Planetary Computer STAC API.”
 - A script with code instructing how to access ABI data from blob storage on Azure.



Visualize with Python: Custom Packages

- The prior Python examples employ widely-used packages for processing any environmental data. However, by using more specialized Python packages, it is possible to process and visualize GOES-R imagery with fewer lines of code. Follow the respective installation guides and documentation.
- [Satpy](#) - a Python package for reading, writing and manipulating satellite imagery by the PyTroll Team.
 - Satpy processes GOES-R data using specialized readers: *abi_l1b*, *abi_l2_nc*, and *glm_l2*. Satpy's capabilities include generating composites, extracting measured values, resampling to a uniform grid, slicing and subsetting scenes.
 - See [Satpy's "Quickstart" webpage](#) for an overview, and "[Reading Data with the Scene](#)" for an ABI L1b example.
 - [Script #5](#): "GOES-16 Mosaic"- Pull files from AWS, or load local files, then join all 16 bands with ImageMagick.
 - [Script #6](#): "GOES-16 ABI - True Color Animation" - Combine single ABI timesteps to create an mp4 video file.

Script #7 changes based on a 7/11/25 review:

```
When calling G = GOES().latest(),  
specify a currently operational satellite,  
e.g., G =  
GOES(satellite=19).latest()
```

```
When calling the nearesttime  
function, change the format to  
GOES(satellite=16,  
domain="F").nearesttime("2021  
-01-01 18:00")
```

```
Comment out any functions from  
"Carpenter_Workshop"
```

- [GOES-2-go](#) - a Python package created by Brian Blaylock, Ph.D.
 - Downloads and reads ABI and GLM data from AWS. A user may retrieve the latest GOES-R data, select images from a specific time and date, or choose a series of images from a date range.
 - Creates multispectral RGB composites, based on recipes from [RAMMB/CIRA](#), such as True Color, Natural Color, Air Mass, and Day Cloud Phase Distinction. See [Script #7](#): "Demo RGB Recipes".
- [goespy](#) - a Python package created by Steven Pestana and Paulo Alexandre Mello.
 - Simplifies downloading from AWS with two functions: *ABI_Downloader* and *GLM_Downloader*.
 - [Script #8](#): "Downloading and plotting GOES-R imagery with goespy and xarray" - Demonstrates goespy utility.

Script #6 changes based on a 7/11/25 review - in the first cell:

```
Add demo to the  
satpy imports:  
from satpy  
import Scene,  
demo
```

```
Adjust the  
BASE_DIR path to a  
local directory, then  
add the line  
satpy.demo.abi_  
l1b.get_hurrica  
ne_florence_abi  
(base_dir=BASE_  
DIR)
```

Visualize with Earth Engine

- [Google Earth Engine](#) (GEE) is a cloud-based platform for geospatial data analysis. The following list summarizes the advantages of imagery processing using GEE:
 - The GOES-R imagery stored in the GEE Data Catalog is already georeferenced. Therefore, it only requires a few lines of script to display an image accurately (Figure 8).
 - GEE provides advanced image processing tools designed for satellite imagery analysis, offered through several APIs.
 - The GEE Data Catalog stores many datasets, including imagery from MODIS, Landsat, and Sentinel, which are updated daily. Synthesizing dissimilar satellite products is possible in GEE.
 - GEE takes advantage of Google's infrastructure to execute high speed parallel processing. This is helpful for handling large collections of images, without using local computing resources.
 - For an introduction to GEE, check out [the GEE Community Tutorials](#).
- Be aware of the following:
 - To use this platform, it is necessary to [apply for a free GEE account](#), which may take a day or two to get approved.
 - [Two GOES-R L2+ products are currently available in the GEE Data Catalog](#): Multi-band Cloud and Moisture Imagery (MCM) and Fire Detection and Characterization (FDC). Sample scripts are available in the dataset descriptions.




Figure 8. GOES-16 CONUS False Color image on June 12th, 2020, visualized using GEE.

Technique: Radiance to Reflectance

- When using the ABI L1b Radiance product, it may be beneficial to convert from the unit radiance, which takes into account the solar irradiance and Earth-Sun distance. For the reflective bands (1-6), convert to reflectance. For the emissive bands (7-16), convert to brightness temperature (Kelvin).
- Reflectance conversion equation: $\text{reflectance } \rho_{f_v} = \text{kappa factor } \kappa * \text{radiance } L_v$
 - The kappa factor is included in the metadata for every L1b file, stored as the variable 'kappa0'
 - For more information on the reflectance conversion, or how to convert to brightness temperature, see the PUG Volume 3: L1b Products, found on the [OSPO GOES-R Documents page](#) (Revision 2.3, pages 27-28).
- Script #9: “True-Color Image: GOES-R ABI L1b Radiances” by Danielle Losos.
 - This Jupyter Notebook demonstrates several useful techniques for processing and visualizing the L1b Radiance product to create a georeferenced RGB True-Color image.
 - Compare to the [Script #3](#) methodology for creating true color images from the L2+ MCMI product.
 - See [Slide 24](#) for background on using multiple bands to generate true-color composite images.
 - Alternative approach for converting radiance to reflectance values is documented in the tutorial: [“Jupyter Notebook for Working with GOES-16 Data”](#) by Opens Commons Consortium (OCC). However, this script errs in using constant values to convert between radiance and reflectance. It is recommended to always use the coefficients embedded in that particular image’s metadata for conversion to reflectance or brightness temperature.
 - A section of the OCC script shows how to visualize Geostationary Lightning Mapper (GLM) point data.

```
Script #9 changes based on a
7/1/25 review:
To avoid an error in cell
In [15] add the following:
import warnings
np.warnings = warnings
```



Technique: Generating Composites

- "Band stacking" is the process of overlaying certain bands to create composite imagery. Read "[Satellite imagery RGB: Adding Value, Saving Time](#)" for background on how composites aid weather forecasting.
- A popular composite image type is an RGB True Color, which uses red, green, and blue wavelength imagery to mimic the appearance of the Earth to the human eye.
 - ABI has blue and red visible channels (bands 1 and 2), but not a green visible channel. The near infrared (NIR) channel (band 3) can serve as a proxy for the green band. [Script #3](#) shows how to create an RGB True Color image from bands 1, 2, and 3 of the MCM1 product. MCM1 is a single file with all 16 bands resampled to 2 km resolution.
 - Using three separate CMI files for the three bands is less data intensive and achieves a higher resolution. However, band 2 has 0.5 km resolution while bands 1 and 3 have 1 km resolution, which makes multi-band operations difficult. Thus, it is necessary to resample band 2. [Script #9](#) demonstrates how to resample using a Rebin function.
- Other ABI RGB composites do not simulate true color, but instead highlight various atmospheric and surface features, such as the [Air Mass RGB](#), [Day Cloud Phase Distinction RGB](#), and [Dust RGB](#).
 - Satpy and GOES-2-Go packages automate the generation of common ABI composites.
- Another common composite is [NDVI](#) (normalized difference vegetation index). This index determines vegetative health by finding the difference between a pixel's reflectance in red and NIR wavelengths.
 - [Script #10](#): Shows how to easily calculate NDVI of a Landsat image in Python using the module Earthpy. To calculate NDVI using ABI, substitute band 2 as the red band and band 3 as the NIR band.

Part 4.




How Can I Process the Data Using GIS?



Make GOES-R Series Data GIS-compatible

- The native format for GOES-R products is netCDF files (see [Slide 30](#)) which are not always compatible with GIS. To process products using GIS software, it may be necessary to change the data format.
- Raster and vector models are the two primary ways to represent geospatial data for visualization in GIS.



Source:
[Saylor Academy](#)

Figure 9. This graphic shows how point/line/polygon vectors and raster grids model surfaces differently.

- Raster Models: grids of rectangular pixels, with each pixel assigned a value that corresponds to the measured environmental feature. To learn more, read "[What is raster data?](#)".
 - Since GOES-R netCDF files store variables in multidimensional arrays, most products can be converted easily to raster file formats ([Slide 27](#)).
 - Over a hundred raster file types are supported by the open-source platform QGIS. Common formats for storing satellite imagery include TIFF, IMG, GRID, JPEG, JP2, BMP, GIF, PNG, BIL/BIP/BSQ, and DAT.
- Vector Models: points, lines, or polygons used to explicitly delineate the boundaries of environmental features. Each geometry (a point/line/polygon) may have associated attributes which describe the characteristics of the feature. For more information, read "[Vector Data](#)".
 - GOES-R netCDF multidimensional arrays cannot be modeled by vectors without transforming the data.
 - Shapefiles are the most widely used vector format and are compatible with any GIS platform. Other file formats for storing vector data include GeoJSON, KML/KMZ, TIN, and DXF.

Convert NetCDF files to GeoTIFFs

- NetCDF files are compatible with select GIS platforms. [QGIS Version 3.2+](#) can handle some GOES-R products. Within QGIS, go to Add Layer > Add Raster Layer, then select a variable when prompted.
- GeoTIFFs are regarded as the industry standard for georeferenced raster data. The conversion of GOES-R netCDF arrays to single-band GeoTIFF rasters can be accomplished in command line, using GDAL or Geo2Grid (see [Slide 19](#)).
 - Install the package [Geospatial Data Abstraction Library \(GDAL\)](#). Use the command `gdal_translate NETCDF:"Input_file.nc":Variable_name Output_file.tif`
 - For example: `gdal_translate NETCDF:"OR_ABI-L1b-RadC-M6C01_G16_s20210010016176_e20210010018549_c20210010019000.nc":Rad Sample_Output.tif`
- The collection of scripts, "[GOES-R NetCDF to GeoTIFFs](#)" by Danielle Losos, can assist with batch retrieval and conversion. These scripts automate the file download by pulling data from a given satellite, product, and date from AWS.




Figure 10. GOES-16 L2+ CMI clipped to the Minnesota border, converted to a GeoTIFF, and visualized in QGIS using Script #13.


Script #12 changes based on a 7/3/25 review: In cell In [9] - change `xr.open_rasterio` to `rioxarray.open_rasterio`, add `tiled="NO"` argument to `netCDF_file.rio.to_raster`

- [Script #11](#): "L1b Channel Loop" - Loops through the 16 netCDF files contained in an ABI L1b Radiance product, and converts each to a GeoTIFF.
- [Script #12](#): "L2 Products" - Visualizes and converts the chosen GOES-R L2+ ABI product variable, e.g., Cloud Top Height, from a netCDF file to a GeoTIFF.
- [Script #13](#): "Clip with Shapefile" - Clips a netCDF file image to a shapefile boundary, and saves the region of interest as a GeoTiff (Fig. 10).

Script #13 changes based on a 7/3/25 review: In cell In [15] - change `xr.open_rasterio` to `rioxarray.open_rasterio`, change `netCDF_file.crs` to `netCDF_file.rio.crs`

Transform Discrete Data to Shapefiles

- Discrete data, also known as thematic or categorical data, is used to represent features that have defined boundaries. Vector models are better suited for discrete than continuous variables.
 - Several ABI L2+ products have discrete variables, including the Clear Sky Mask which provides a binary classification for each pixel: “clear” or “cloudy”; Aerosol Detection; and Cloud Top Phase.
 - Most GOES-R products have a Data Quality Flag (DQF) layer, a discrete variable which classifies each pixel’s usability into categories like “good quality”, “conditionally usable”, or “out of range”.
- [Script #14: “Discrete variable to shapefile”](#) by Danielle Losos.
 - This script transforms ABI products to shapefiles by polygonizing gridded data. A vector polygon is created for each connected region of pixels in the array sharing a common value.
 - For example, the Clear Sky Mask can be polygonized to a vector layer where each polygon has an attribute equal to 0 for cloud-free or 1 for cloud-covered (Fig. 11).



0 ABI L2+ Clear Sky Mask 1




Figure 11. Clear Sky Mask vector layer (left image) is used to mask an RGB-false color raster in QGIS (right image).

Script #14 changes based on a 7/3/25 review:

In the first cell, add `import rioxarray`

In cell In [9] -

change `xr.open_rasterio` to `rioxarray.open_rasterio`,

change `netCDF_file.crs` to `netCDF_file.rio.crs`,

change `Projection.ImportFromProj4(goesR_crs)` to

`Projection.ImportFromProj4(goesR_crs.to_proj4())`

Part 5.



Frequently Asked Questions





How are GOES-R Series files formatted?

- GOES-R Series product files use the netCDF-4 format, a general-purpose scientific data file format.
- More details on netCDF files:
 - GOES-R Series products may have any of the following types of attributes
 - **title** : a global attribute that is a character array providing a succinct description of what is in the dataset.
 - **Conventions** : a global attribute that is a character array for the name of the conventions followed by the product.
 - **long_name** : long descriptive name for each variable.
 - **_FillValue** : a scalar value that identifies missing data.
 - **Valid_range** : a delimited vector of two numbers specifying the minimum and maximum valid values for the variable to which it is attached.
 - **scale_factor** and **add_offset** : these attributes are used together to provide simple data compression to store floating-point data as small integers in a product data file.
 - **units** : a character string that specifies the units used for the variable's data.
 - A netCDF file also includes the dimensions that are used to size dimensional variables.
 - For a quick look at the contents of a netCDF file, use the `ncdump` command line tool from [Unidata's netCDF package](#).
 - The netCDF ABI L1b/L2+ gridded product files are compressed to reduce the file size. For information on how to unpack data, see [Slide 31](#). For more details on netCDF files, refer to the PUG Volume 1: Main, found on the [OSPO GOES-R Documents page](#) (Revision 2.3, pages 8-14).
- Other file formats:
 - Flexible Image Transport System (FITS) format is used alongside netCDF for the SUVI Solar EUV Imagery product.
 - FITs is the primary format used for the CCOR data products, with their auxiliary files in netCDF format.
 - Unix text file format is used in a small subset of the Level 1b and 2+ semi-static source data files.
 - Hierarchical Data Format (HDF) is used for several Level 1b semi-static source data files.



How do I pre-process GOES-R Series data?

- GOES-R Series data are packed in compressed files, which must be “unpacked” for integer analysis. Images can be visualized without unpacking, since scaled integers are still proportional to each other.
- What is packed data?
 - In order to minimize file size, many of the ABI L1b/L2+ products use 16-bit **scaled** integers for physical data quantities rather than 32-bit floating point values.
 - To convert data back to the actual value associated with the physical quantity, the user must multiply by a scale factor and add an offset to the 16-bit scaled integer.
- How do I unpack data?
 - To unpack the data, apply the attributes using the following equation:
$$\text{unpacked_value} = \text{packed_value} * \text{scale_factor} + \text{add_offset}$$
 - The variables ‘scale_factor’ and ‘add_offset’ are included in the metadata for each file. The scale factor is calculated with the formula (Max Value - Min Value)/65530, and the offset is the product’s expected minimum value.
 - Here is a sample conversion for band 8: $\text{Brightness temperature(Kelvin)} = (2,305 * 0.04225) + 138.05 = 235.44 \text{ K}$
 - Before unpacking, check to ensure that the program in use is not automatically applying the scale factor and offset.
- How do I remove data that is fill, out of range, or missing, from the image?
 - These faulty data types can be masked using data quality flags (DQFs), convention attributes which associate integer values with a categorization of the data quality.
 - For more information on packed data or DQFs, see the PUG Volume 1: Main, found on the [OSPO GOES-R Documents page](#) (Revision 2.3, pages 21 or 16, respectively).

How are ABI L1b products georeferenced?

- ABI uses a fixed grid whose native coordinate values are the East/West scanning angle and North/South elevation angle in units of radians relative to the location of the satellite.
 - The ABI fixed grid is a projection based on the viewing perspective of the idealized location of a satellite in geosynchronous orbit. This allows the same data points in every product to be at the same location on Earth.
 - The fixed grid is rectified to an ellipsoid defined by the Geodetic Reference System 1980 (GRS80) earth model.
 - For all ABI gridded L1b/L2+ products, the names of the coordinate variables are x (E/W scanning angle) and y (N/S elevation angle) (Figure 13).
 - Additionally, there is a “grid_mapping” attribute which is attached to data variables whose values are associated with specific earth locations.
 - For ABI L1b/L2+ gridded products, the grid mapping attribute specifies the projection (GRS80), the lat/lon origin of the projection, and a parameter that identifies the ABI scanning pattern.
- Grid mappings coupled with the coordinate variables provide the means to determine the latitude and longitude of a data point.
 - See the PUG Volume 3: L1b Products, found on the [OSPO GOES-R Documents page](#) (Revision 2.3, pages 8-26), for an in-depth explanation.




Figure 13. Every data point is plotted on the ABI fixed grid as a distance from the (0,0) origin, in units x (East to West Scanning Angle) and y (North to South Elevation Angle).



Why are data not available on a given date?

- Cloud-based platforms have select L2+ Product availability.
 - Once a GOES-R Series product has reached provisional maturity status, cloud-based platforms may release the product.
 - When there is demonstrated user interest in a product, vendors may backfill their archive to include the dates requested. This may include beta maturity levels of the product.
 - If there is no demonstrated user interest in the product, vendors may choose not to release the entire available collection. For this reason, collection length may vary between products on cloud-based platforms.
- To retrieve pre-provisional GOES-R Series products, search on NOAA CLASS to access the entire archive of products made available to users. CLASS stores products with beta, provisional, and full maturity status.
- Occasionally, there is missing GOES-R Series data on select dates and times.
 - To investigate whether data is missing due to issues with data reception, check out the [SSEC's Monitoring Web Page](#). Follow these instructions for tips on how to navigate this website:
 1. First, select the GOES-R Series satellite of interest, then choose a date of interest at the top of the page.
 2. Each row represents the data quality of full disk, CONUS, and mesoscale images for every hour of the day.
 3. The "DB: SSEC" section reports missing data due to GOES Rebroadcast (GRB) reception issues.
KEY- **light green**: 100% received, **yellow**: 96 -100% received, **light red**: < 96% received, **dark red**: missing
 4. The "PDA via STAR" section reports missing data due to errors with Product Distribution and Access (PDA).
KEY (same as above with two more categories) - **blue** & **orange**: data available from PDA but missing or late
 - Also, check out [OSPO's Satellite Alert Messages](#) to see if there were any outages reported on that day



Other Questions and Contact

Other Questions

- For unanswered questions, try browsing the [GOES-R Series FAQ Page](#) or [NCEI's GOES-R Series Page](#).
- Additional resources are available on the [Office of Satellite and Product Operations \(OSPO\) GOES-R documents page](#), including links to the most recent Product User's Guides (Volumes 1-5) and the Product Algorithm Theoretical Basis Documents (ATBDs). The [GOES-R Series Documents page](#) also contains links to many helpful GOES-R documents.
- Useful links for background information: [Mission Overview](#), [GOES History](#), [GOES-R Data Book](#) (technical overview of satellite and ground systems)

Contact

- Get in touch by emailing the Satellite Products and Services Division (SPSD) User Services at SPSD.UserServices@noaa.gov. Please share your questions, concerns, or suggestions for how this GOES-R beginner's guide could be more helpful.
- Show us how you are using GOES-R data in your own work by using [#goesrguide](#) when you post on social media!



Appendix A: Acronym List

ABI - Advanced Baseline Imager
AIRS - Archive Information Request System
ATBD - Algorithm Theoretical Basis Documents
AWIPS - Advanced Weather Interactive Processing System
AWS - Amazon Web Service
CAVE - Common AWIPS Visualization Environment
CCOR - Compact Coronagraph
CIMSS - Cooperative Institute for Meteorological Satellite Studies
CIRA - Cooperative Institute for Research in the Atmosphere
CLASS - Comprehensive Large Array-data Stewardship System
CMI - Cloud and Moisture Imagery
CONUS - Continental U.S.
CSSP - Community Satellite Processing Package
DQF - Data Quality Flag
EHIS - Energetic Heavy Ion Sensor
EUVS - Extreme Ultraviolet Sensor
EXIS - Extreme Ultraviolet and X-ray Irradiance Sensors
FDC - Fire Detection and Characterization

FITS - Flexible Image Transport System
FTP - File Transfer Protocol
GDAL - Geospatial Data Abstraction Library
GEE - Google Earth Engine
GIS - Geographic Information System
GLM - Geostationary Lightning Mapper
GOES - Geostationary Operational Environmental Satellite
GRB - GOES Rebroadcast
GRS80 - Geodetic Reference System 1980
HDF - Hierarchical Data Format
L0/L1b/L2+ - Level 0/ Level 1b/ Level 2+
MAG - Magnetometer
MCMI - Multi-band Cloud and Moisture Imagery
MESO 1/2 - Mesoscale Domain Sector 1/2
MPS-HI/LO - Magnetospheric Particle Sensor - High/Low
NASA - National Aeronautics and Space Administration
NCEI - National Centers for Environmental Information
NDVI - Normalized Difference Vegetation Index
NetCDF - Network Common Data Format

NIR - Near-infrared
NOAA - National Oceanic and Atmospheric Administration
NWS - National Weather Service
OCC - Opens Commons Consortium
OSPO - Office of Satellite and Product Operations
PACUS - Pacific U.S.
PDA - Product Distribution and Access
PUG - Product User Guide
RAMMB - Regional and Mesoscale Meteorology Branch
RGB - Red-Green-Blue
SEISS - Space Environment In-Situ Suite
SGPS - Solar and Galactic Proton Sensor
SPoRT - Short-term Prediction Research and Transition Center
SPSD - Satellite Products and Services Division
SSEC - Space Science and Engineering Center
SUVI - Solar Ultraviolet Imager
SWPC - Space Weather Prediction Center
WFO - Weather Forecast Office

Additional acronyms defined on the [GOES-R Series Acronym Page](#)



Appendix B: Deprecated Script Examples

- Slides in this appendix were included in a previous version of the GOES-R Beginner's Guide, but the script examples include older Python functions that are now deprecated
- We are continuing to include these older slides in case the functional parts of the scripts provide enough information for a user to start using it and possibly find a workaround on their own

Transform Continuous Data to Shapefiles

- Continuous data represents variables that change progressively across surfaces. Most GOES-R products contain continuous variables, like radiance, reflectance, temperature, or pressure.
 - Continuous variables are not modeled well by vector data. Transforming satellite imagery to vectors requires generalization of data into categories (i.e., discrete intervals) which results in a loss of detail and accuracy.
 - However, certain geospatial analytic operations can only be performed on vector data.
- Script #B-1: “Continuous variable to contour to shapefile” by Danielle Losos.
 - This script converts ABI products to shapefiles by first transforming the continuous variable into a contour plot of n discrete intervals. Next, the concentric rings of the contour plot are converted into shapefile polygons.
 - As the number of contour intervals, n , increases, the accuracy of the vector model improves (Figure B-1). The trade off for a higher n is a longer processing time and ultimately a larger file size.

Script #B-1 changes:
 Perform similar changes as described in the Script #14 changes
 The actual shapefile creation in cells IN 10-12 include deprecated functions, and the direct replacements produce unrealistic results




Figure B-1. GOES-16 CONUS L2+ Land Surface Temperature visualized as a contour plot with 5, 15, and 50 discrete intervals.